



IMMUNISATION SCHEDULE

The table below shows the timetable for immunisations but your GP or health visitor should remind you when the next one is due. Try to stick to the immunisation schedule that your GP gives you to ensure that your child is protected as soon as possible.

0-5 months

2 Months

Diphtheria, tetanus, pertussis (whooping cough), polio, and Hib (DTaP/IPV/Hib)

Combined Injection

You may see a small lump appear where the injection was given.*

Immunisation due:

Pneumococcal conjugate vaccine (PCV)

Separate Injection

Possible redness and swelling where the injection was given. Mild fever.*

Immunisation due:

- What is Given
- How it is Given
- Possible side effects



3 Months

Diphtheria, tetanus, pertussis (whooping cough), polio and Hib (DTaP/IPV/Hib)

Combined Injection

Post - immunisation fever within 12 – 24 hours.*

Immunisation due:

MEN C vaccine (Meningitis C injection)

Separate Injection

Possible redness and swelling where the injection was given. Mild fever.*

Immunisation due:

4 Months

Diphtheria, tetanus, pertussis (whooping cough), polio and Hib (DTaP/IPV/Hib)

Combined Injection

Post - immunisation fever within 12 – 24 hours.*

Immunisation due:

MEN C vaccine (Meningitis C injection)

Separate Injection

Possible redness and swelling where the injection was given. Mild fever.*

Immunisation due:

Pneumococcal conjugate vaccine (PCV)

Separate Injection

Possible redness and swelling where the injection was given. Mild fever.*

Immunisation due:

1-2 Years

12 Months

Hib and Meningitis C (Hib/MenC) booster

Combined Injection

Post - immunisation fever within 12 – 24 hours.*

Immunisation due:

Approx 13 Months

MMR vaccine (Measles, Mumps and Rubella)

Combined Injection

Post - immunisation fever within 12 – 24 hours.*

Immunisation due:

Pneumococcal conjugate vaccine (PCV)

Separate Injection

Possible redness and swelling where the injection was given. Mild fever.*

Immunisation due:

3-5 Years

40 months

Diphtheria, tetanus, pertussis (whooping cough) and polio (DTaP/IPV)

Combined Injection

Post - immunisation fever within 12 – 24 hours.*

Immunisation due:

MMR vaccine (Measles, Mumps and Rubella)

All in one Injection

Possible fever and measles - like rash.*

Immunisation due:

6+

13-18 years

Diphtheria, tetanus, and polio (Td/IPV)

Combined Injection

You may see a small lump appear where the injection was given.*

Immunisation due:

Girls 12-13

Human Papillomavirus (HPV)

3 Injections over 6 months

Possible swelling, redness and pain where the injection was given.*

Immunisation due:



* This is one possible side effect; for more side effects visit www.immunisation.nhs.uk

Always read the label. Calpol Infant Suspension contains paracetamol. For pain and fever relief